

Foveal Crowding: Pilot results

Mapping Acuity and Crowding Effects in the Fovea

Janis Intoy, Radhika Dhanak

APLAB

August 4, 2017

Acuity across the Retina

Acuity is measured as minimum angle of resolution (MAR) or in log units as LogMAR.





Figure: Anderson et al. (1991) and Jacobs (1979)

Acuity in the Foveola??




Figure: Marcos & Navarro (1997)




Figure: Rossi & Roorda (2010)

How are the effects of crowding measured?

Approach 1: Quantify impact of flanker(s) on task performance using fixed target and flanker sizes and varying flanker distance or position. (Bouma, 1970; Nandy & Tjan, 2012)




Figure: Bouma (1970) and Nandy & Tjan (2012)

How are the effects of crowding measured?

Approach 2: Quantify impact of flanker(s) with threshold estimations for varying target and flanker sizes, flanker distances, eccentricities,... (Pelli et al, 2004; Chung et al, 2001; Levi et al., 2002)




Figure: Pelli et al. (2007)

How are the effects of crowding measured?

Approach 3: Quantify impact of flanker(s) by measuring flanked and unflanked acuity where the flanker distance is a multiple of the target size (Levi, Song, & Pelli, 2007; Jeon et al, 2010; Norgett & Siderov, 2014).

•
R
Identify one
isolated letter.

•
HRO
Z
Identify one
flanked letter.




Figure: Levi, Song, & Pelli (2007)

What about foveal crowding?

“[acuity] is unaffected by crowding if done foveally, where critical spacing is only a few minutes of arc” (Pelli & Tillman, 2008)

“there is a question about whether genuine crowding actually occurs in the fovea” (Levi, 2008)

What about foveal crowding?

- technical difficulties in presenting stimulus at different foveal eccentricities
- crowding and masking may be confused within the fovea because of optical blur
- more general masking effects, not necessarily crowding (for: Levi et al., 2002; Hariharan et al., 2005; against: Danilova & Bondarko, 2006)
- physics, not physiology (Hess et al., 2000; Liu & Arditi, 2000; Liu 2001; Chung & Tjan, 2007)





Figure: Liu (2000); Danilova & Bondarko (2006)

Objectives

- Map acuity within the foveola
 - asymmetries in nasal vs temporal foveal vision?
 - individual variability?
- Measure the effects of crowding within the foveola

(very rough) schematic diagram of display setup

We need high spatial and temporal resolution to measure acuity under stabilized conditions.



- eye to display = 4.1m
- pixel angle = 0.236 arcmin and 200Hz refresh rate

Experiment Design: Overview

We want to measure visual acuity in uncrowded and crowded conditions.


- Eccentricities tested: 0, 5, 15, 25, and 60 arcmin¹
- 10 conditions (uncrowded/crowded x eccentricity) tested in individual blocks of about 100 trials in pseudorandom order
- Task: Report orientation of tumbling-E optotype (left or right, 2-AFC)
- PEST algorithm changed size of optotypes to target 75% performance in discrimination task
- Stimuli were fully stabilized to maintain fixed eccentricity from the center of gaze

Recalibration every trial.

1.5 second fixation every 15 trials.

¹only temporal eccentricities tested in pilot


Experiment Design: Stimuli



Examples of uncrowded and crowded stimuli.


- All tumbling-Es were horizontally facing (left or right).
- Edge-to-edge distance between target and flankers is half the size of the target.

Experiment Design: Display Dimensions



Note that central fixation point is shown only for eccentricities $> 5'$.

Experiment Design: Trial Flow



Data Collection: Overview

- $N = 2$
- ideally we would only include drift trials but we may include MS trials if yields are low
- Today we will mostly look at data from the naive subject (drift only trials)



Data Collection: Naive Subject

Full data completed for one naive subject (temporal eccentricities only).
(3 training sessions, 6 data collection sessions)


	Total	Valid	MS	S	NT/B	Yield
All	2004	1223	132	646	3	67.61%
Uncrowded	1018	634	72	310	2	69.35%
Crowded	986	589	60	336	1	65.82%
Fixation	125	0	30	45	50	24.00%

Ecc.	Uncrowded						Crowded					
	Total	Valid	MS	S	NT/B	Yield	Total	Valid	MS	S	NT/B	Yield
0	143	96	15	31	1	111: 77.62%	155	107	24	24	0	131: 84.52%
5	160	125	13	22	0	138: 86.25%	254	162	25	66	1	187: 73.62%
15	184	143	8	33	0	151: 82.07%	159	126	2	31	0	128: 80.50%
25	253	129	12	112	0	141: 55.73%	170	96	5	69	0	101: 59.41%
60	278	141	24	112	1	165: 59.35%	248	98	4	146	0	102: 41.13%


Saccade and Microsaccade rates



PDF of gaze positions - part 1



PDF of gaze positions - part 2



Psychometric Function fitting

$$P(x) = \gamma + S(x) \cdot (1 - (\lambda + \gamma))$$

x = stimulus size

$\gamma = 0.5$

$\lambda = 0$

psychometric function (MAR)
(guess rate)
(lapse rate)

- Cumulative normal distribution

$$S(x) = \frac{1}{2} \left[1 + \operatorname{erf} \left(\frac{x - \mu}{\sigma\sqrt{2}} \right) \right]$$


- Cumulative Weibull distribution²

$$S(x) = \begin{cases} 1 - e^{-(x/\alpha)^\beta}, & x \geq 0 \\ 0, & x < 0 \end{cases}$$




²Used to measure acuity thresholds by Williams & Coletta (1987), Lalor et al. (2016), Marcos & Navarro (1997), Jeon et al. (2010) and others.

Eccentricity = 0 arcmin

Normal




Weibull






Eccentricity = 5 arcmin

Normal




Weibull






Eccentricity = 15 arcmin

Normal




Weibull






Eccentricity = 25 arcmin


Normal




Weibull



Eccentricity = 60 arcmin



MAR Acuity by Eccentricity (Naive subject)



MAR Acuity by Eccentricity: Threshold Values

Drift and MS

Ecc	Cumulative Normal distribution			
	Uncrowded		Crowded	
	Thresh	Boot mn±sd	Thresh	Boot mn±sd
0	1.33	1.33±0.15	1.22	1.22±0.12
5	1.59	1.58±0.11	1.09	1.09±0.12
15	1.49	1.49±0.21	1.55	1.55±0.14
25	1.55	1.55±0.11	1.87	1.85±0.16
60	1.26	1.27±0.11	4.00	3.99±0.35

Ecc	Cumulative Weibull distribution			
	Uncrowded		Crowded	
	Thresh	Boot mn±sd	Thresh	Boot mn±sd
0	1.37	1.37±0.15	1.24	1.23±0.15
5	1.63	1.61±0.14	1.06	1.07±0.16
15	1.52	1.52±0.26	1.59	1.58±0.14
25	1.59	1.58±0.11	1.89	1.87±0.16
60	1.23	1.25±0.14	4.07	4.04±0.36

Drift Only

Ecc	Cumulative Normal distribution			
	Uncrowded		Crowded	
	Thresh	Boot mn±sd	Thresh	Boot mn±sd
0	1.51	1.50±0.14	1.16	1.15±0.15
5	1.60	1.59±0.11	1.08	1.07±0.13
15	1.50	1.50±0.24	1.55	1.54±0.15
25	1.55	1.55±0.11	1.88	1.86±0.15
60	1.29	1.29±0.12	4.07	4.09±0.38

Ecc	Cumulative Weibull distribution			
	Uncrowded		Crowded	
	Thresh	Boot mn±sd	Thresh	Boot mn±sd
0	1.52	1.51±0.15	1.16	1.17±0.18
5	1.63	1.62±0.13	1.06	1.05±0.10
15	1.53	1.55±0.28	1.59	1.58±0.16
25	1.58	1.58±0.12	1.90	1.89±0.16
60	1.27	1.29±0.14	4.13	4.10±0.34

*1000 bootstrap iterations

Data from Experienced Subject

- Eccentricities: 0, 5, and 60arcmin
- Analyzed drift and MS trials

Data Collection: Experienced Subject

Partial data collected for one experienced subject.
(3 data collection sessions)

	Total	Valid	MS	S	NT/B	Yield
All	1062	513	353	188	8	81.54%
Uncrowded	528	241	234	50	3	89.96%
Crowded	534	272	119	138	5	73.22%
Fixation	86	1	76	8	1	89.53%

Ecc.	Uncrowded						Crowded					
	Total	Valid	MS	S	NT/B	Yield	Total	Valid	MS	S	NT/B	Yield
0	214	93	116	4	1	209: 97.66%	135	94	38	3	0	132: 97.78%
5	118	48	59	10	1	107: 90.68%	127	85	25	17	0	110: 86.61%
15	41	20	16	5	0	36: 87.80%	35	11	13	10	1	24: 68.57%
25	30	9	11	9	1	20: 66.67%	21	3	3	14	1	6: 28.57%
60	125	71	32	22	0	103: 82.40%	216	79	40	94	3	119: 55.09%

MAR Acuity by Eccentricity: Threshold Values

Drift and MS

Ecc	Cumulative Normal distribution				Ecc	Cumulative Weibull distribution			
	Uncrowded		Crowded			Uncrowded		Crowded	
	Thresh	Boot mn±sd	Thresh	Boot mn±sd	Thresh	Boot mn±sd	Thresh	Boot mn±sd	
0	1.11	1.10±0.12	1.22	1.21±0.08	0	1.13	1.12±0.11	1.23	1.23±0.08
5	1.05	1.04±0.15	1.19	1.19±0.09	5	1.06	1.06±0.15	1.20	1.20±0.09
15	1.32	1.31±0.11	1.42	1.40±0.09	15	1.33	1.33±0.11	1.43	1.42±0.10
25	0.98	1.14±0.16	NaN	1.73±0.08	25	0.97	1.16±0.16	NaN	1.74±0.08
60	1.22	1.20±0.22	2.13	2.11±0.22	60	1.24	1.23±0.23	2.16	2.13±0.24

*1000 bootstrap iterations

MAR Acuity by Eccentricity (Experienced subject)

